

0.25% | 0.5%

Bupivacaine Hydrochloride Injection, USP



AP Rated

Preservative-Free

Not Made with Natural Rubber Latex

WARNING: RISK OF CARDIAC ARREST WITH USE OF BUPIVACAINE HYDROCHLORIDE INJECTION IN OBSTETRICAL ANESTHESIA

See Full prescribing information for complete boxed warning.

There have been reports of cardiac arrest with difficult resuscitation or death during use of bupivacaine hydrochloride injection for epidural anesthesia in obstetrical patients. In most cases, this has followed use of the 0.75% (7.5 mg/mL) concentration. Resuscitation has been difficult or impossible despite apparently adequate preparation and appropriate management. Cardiac arrest has occurred after convulsions resulting from systemic toxicity, presumably following unintentional intravascular injection. The 0.75% (7.5 mg/mL) concentration of bupivacaine hydrochloride injection is not recommended for obstetrical anesthesia and should be reserved for surgical procedures where a high degree of muscle relaxation and prolonged effect are necessary (5.1).

NDC	Strength	Package Quantity	Wholesaler Information	
72266-263-10	25 mg/10 mL (2.5 mg/mL)	10 Single-Dose Vials	Cardinal Cencora McKesson	6106629 10307348 3082567
72266-264-10	50 mg/10 mL (5 mg/mL)	10 Single-Dose Vials	Cardinal Cencora McKesson	6106637 10307188 3082559



SCAN TO VIEW
OUR PRODUCTS

To order, call (855) 282-4882

Fosun Pharma USA Inc.
104 Carnegie Center, Suite 204 Princeton, NJ 08540
fosunpharmausa.com